ONDERSTEPOORT JOURNAL OF VETERINARY RESEARCH

Volume 70 • Numbers 1–4 • 2003

Author index

AYLING, R.

Comparison of complement fixation test, immunoblotting, indirect ELISA, and competitive ELISA for detecting antibodies to *Mycoplasma mycoides* subspecies *mycoides* small colony (SC) in naturally infected cattle from the 1995 outbreak in Botswana—C. MAROBELA-RABOROKGWE, R. NICHOLAS, R. AYLING and J.B. BASHIRUDIN ... 21–27

BAIPOLEDI, E.K.

Isolation of *Toxoplasma gondii* from goats with a history of reproductive disorders and the prevalence of *Toxoplasma* and chlamydial antibodies—S.P. SHARMA, E.K. BAIPOLEDI, J.F.C. NYANGE and L. TLAGAE 65–68

BASHIRUDDIN, J.B.

Comparison of complement fixation test, immunoblotting, indirect ELISA, and competitive ELISA for detecting antibodies to *Mycoplasma mycoides* subspecies *mycoides* small colony (SC) in naturally infected cattle from the 1995 outbreak in Botswana—C. MAROBELA-RABOROKGWE, R. NICHOLAS, R. AYLING and J.B. BASHIRUDIN ... 21–27

BASSON, K.M.

Conditioned feed aversion as a means to prevent tulp (*Homeria pallida*) poisoning in cattle—L.D. SNYMAN, R.A. SCHULTZ, J.P.J. JOUBERT, K.M. BASSON and L. LABUSCHAGNE 43–48

BAULE, C.

Molecular analysis of bovine viral diarrhoea virus isolates from South Africa—N. KABONGO, C. BAULE and M. VAN VUUREN 273–279

BEAUCOURNU, J-C.

Fleas of elephant shrews (Mammalia, Macroscelididae), and a new host and locality record for *Macroscelidopsylla albertyni* De Meillon & Marcus, 1958 (Siphonaptera, Chimaeropsyllidae)—J-C. BEAUCOURNU, I.G. HORAK and L.J. FOURIE 251–253

BEKKER, P.A.

The heart remembers: observations of cardiac memory in the Dorper sheep heart—J. KER, E.C. WEBB, J.A. KER and P.A. BEKKER 299–305

BELOZEROV, V.N.

Population structure of *Argas arboreus* (Acari: Argasidae) ticks associated with seasonally abandoned mixed heronries, dominated by cattle egrets (*Bubulcus ibis*), in South Africa—V.N. BELOZEROV, D.J. VAN NIEKERK and H.J.B. BUTLER 325–330

BHEBHE, E.

Effect of host age in the distribution of adult *Trichinella zimbabwensis* in the small intestines of golden hamsters (*Mesocricetus auratus*) and Balb C mice—S. MUKARATIRWA, E. NKULUNGO, ELIZABETH MATENGA and E. BHEBHE 169–173

BOOMKER, J.

Studies on the genus *Setaria* Viborg, 1795 in South Africa. II. *Setaria scalprum* (Von Linstow, 1908) and *Setaria saegeri* (Le Van Hoa, 1961)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 7–13

Parasites of South African wildlife. XVII. *Ostertagia triquetra* n. sp. (Nematoda: Trichostrongylina) from the grey rhebuck, *Pelea capreolus* (Forster, 1790)—J. BOOMKER and M-C. DURETTE-DESSET 37–41

Parasites of domestic and wild animals in South Africa. XLI. Arthropod parasites of impalas, *Aepyceros melampus* in the Kruger National Park—I.G. HORAK, G.J. GALLIVAN, L.E.O. BRAACK, J. BOOMKER and V. DE VOS ... 131–163

BRAACK, L.E.O.

Parasites of domestic and wild animals in South Africa. XLI. Arthropod parasites of impalas, *Aepyceros melampus* in the Kruger National Park—I.G. HORAK, G.J. GALLIVAN, L.E.O. BRAACK, J. BOOMKER and V. DE VOS ... 131–163

BRAGG, R.R.

Continuous disinfection as a means to control infectious diseases in poultry. Evaluation of a continuous disinfection programme for broilers–R.R. BRAGG and P. PLUMSTEAD 219–229

BROUQUI, P.

First evidence of *Anaplasma platys* in *Rhipicephalus sanguineus* (Acari: Ixodida) collected from dogs in Africa—Y.O. SANOGO, B. DAVOUST, H. INOKUMA, J-L. CAMICAS, P. PAROLA and P. BROUQUI 205–212

BURRIDGE, M.J.

Comparing the detection of exposure to *Ehrlichia ruminantium* infection on a heartwater-endemic farm by the pCS20 polymerase chain reaction assay and an indirect MAP1-B enzyme linked immunosorbent assay—B.H. SIMBI, T.F. PETER, M.J. BURRIDGE and S.M. MAHAN 231–235

BUTLER, H.J.B.

Population structure of *Argas arboreus* (Acari: Argasidae) ticks associated with seasonally abandoned mixed heronries, dominated by cattle egrets (*Bubulcus ibis*), in South Africa—V.N. BELOZEROV, D.J. VAN NIEKERK and H.J.B. BUTLER 325–330

BUYU, G.E.

Use of rapid rural appraisal and cross-sectional studies in the assessment of constraints in small-holder cattle production systems in the western Kenya highlands—O.S. OKUTHE, A. MCLEOD, J.M. OTTE and G.E. BUYU ... 237–242

CAMICAS, J-L.

First evidence of *Anaplasma platys* in *Rhipicephalus sanguineus* (Acari: Ixodida) collected from dogs in Africa—Y.O. SANOGO, B. DAVOUST, H. INOKUMA, J-L. CAMICAS, P. PAROLA and P. BROUQUI 205–212

COHEN, A.S.

Indirect enzyme-linked immunosorbent assay for the detection of antibody against Rift Valley fever virus in domestic and wild ruminant sera—J.T. PAWESKA, S.J. SMITH, I.M. WRIGHT, R. WILLIAMS, A.S. COHEN, A.A. VAN DIJK, A.A. GROBBELAAR, J.E. CROFT, R. SWANEPOEL and G.H. GERDES 49–64

COLEMAN, P.G.

A comparative study on the clinical, parasitological and molecular diagnosis of bovine trypanosomosis in Uganda—J.W. MAGONA, J.S.P. MAYENDE, W. OLAHO-MUKANI, P.G. COLEMAN, N.N. JONSSON, S.C. WELBURN and M.C. EISLER 213–218

CROFT, J.E.

Indirect enzyme-linked immunosorbent assay for the detection of antibody against Rift Valley fever virus in domestic and wild ruminant sera—J.T. PAWESKA, S.J. SMITH, I.M. WRIGHT, R. WILLIAMS, A.S. COHEN, A.A. VAN DIJK, A.A. GROBBELAAR, J.E. CROFT, R. SWANEPOEL and G.H. GERDES 49–64

DAVOUST, B.

First evidence of *Anaplasma platys* in *Rhipicephalus sanguineus* (Acari: Ixodida) collected from dogs in Africa—Y.O. SANOGO, B. DAVOUST, H. INOKUMA, J-L. CAMICAS, P. PAROLA and P. BROUQUI 205–212

DE VOS, V.

Parasites of domestic and wild animals in South Africa. XLI. Arthropod parasites of impalas, *Aepyceros melampus* in the Kruger National Park—I.G. HORAK, G.J. GALLIVAN, L.E.O. BRAACK, J. BOOMKER and V. DE VOS ... 131–163

DU TOIT, J.T.

Seroconversion in captive African wild dogs (*Lycaon pictus*) following administration of a chicken head bait/SAG-2 oral rabies vaccine combination—D.L. KNOBEL, A. LIEBENBERG and J.T. DU TOIT 73–77

DURETTE-DESSET, M-C.

Parasites of South African wildlife. XVII. *Ostertagia triquetra* n. sp. (Nematoda: Trichostrongylina) from the grey rhebuck, *Pelea capreolus* (Forster, 1790)—J. BOOMKER and M-C. DURETTE-DESSET 37–41

EISLER, M.C.

A comparative study on the clinical, parasitological and molecular diagnosis of bovine trypanosomosis in Uganda—J.W. MAGONA, J.S.P. MAYENDE, W. OLAHO-MUKANI, P.G. COLEMAN, N.N. JONSSON, S.C. WELBURN and M.C. EISLER 213–218

ESTRADA-PEÑA, A.

Climate change decreases habitat suitability for some tick species (Acari: Ixodidae) in South Africa—A. ESTRADA-PEÑA 79–93

FOURIE, L.J.

Fleas of elephant shrews (Mammalia, Macroscelididae), and a new host and locality record for *Macroscelidopsylla albertyni* De Meillon & Marcus, 1958 (Siphonaptera, Chimaeropsyllidae)—J-C. BEAUCOURNU, I.G. HORAK and L.J. FOURIE 251–253

GALLIVAN, G.J.

Parasites of domestic and wild animals in South Africa. XLI. Arthropod parasites of impalas, *Aepyceros melampus* in the Kruger National Park—I.G. HORAK, G.J. GALLIVAN, L.E.O. BRAACK, J. BOOMKER and V. DE VOS ... 131–163

GAYO, V.

PCR-based detection of the transovarial transmission of Uruguayan *Babesia bovis* and *Babesia bigemina* vaccine strains—V. GAYO, M. ROMITO, L. NEL, M.A. SOLARI and G.J. VILJOEN 197–204

GERDES, G.H.

Indirect enzyme-linked immunosorbent assay for the detection of antibody against Rift Valley fever virus in domestic and wild ruminant sera—J.T. PAWESKA, S.J. SMITH, I.M. WRIGHT, R. WILLIAMS, A.S. COHEN, A.A. VAN DIJK, A.A. GROBBELAAR, J.E. CROFT, R. SWANEPOEL and G.H. GERDES 49–64

GROBBELAAR, A.A.

Indirect enzyme-linked immunosorbent assay for the detection of antibody against Rift Valley fever virus in domestic and wild ruminant sera—J.T. PAWESKA, S.J. SMITH, I.M. WRIGHT, R. WILLIAMS, A.S. COHEN, A.A. VAN DIJK, A.A. GROBBELAAR, J.E. CROFT, R. SWANEPOEL and G.H. GERDES 49–64

GROENEWALD, H.B.

The morphological characteristics of the antebrachiocarpal joint of the cheetah (*Acinonyx jubatus*)—L.O.C. OHALE and H.B. GROENEWALD 15–20

HARDY, W.D. Jr

Evaluation of an indirect enzyme-linked immunosorbent assay for the detection of feline lentivirus-reactive antibodies in wild felids, employing a puma lentivirus-derived synthetic peptide antigen—M. VAN VUUREN, E. STYLIANIDES (nèe DE KLERK), S.A. KANIA, E.E. ZUCKERMAN and W.D. HARDY Jr 1–6

HORAK, I.G.

Ixodid ticks of Angora and Boer goats, grysbok, common duikers, kudus and scrub hares in Valley Bushveld in the Eastern Cape Province, South Africa—K.M. DE F. MACIVOR and I.G. HORAK 113–120

Parasites of domestic and wild animals in South Africa. XLI. Arthropod parasites of impalas, *Aepyceros melampus* in the Kruger National Park—I.G. HORAK, G.J. GALLIVAN, L.E.O. BRAACK, J. BOOMKER and V. DE VOS ... 131–163

Parasites of domestic and wild animals in South Africa. XLII. Helminths of sheep on four farms in the Eastern Cape Province—I.G. HORAK 175–186

Parasites of domestic and wild animals in South Africa. XLIII. Ixodid ticks of domestic dogs and cats in the Western Cape Province—I.G. HORAK and SONJA MATTHEE 187–195

Fleas of elephant shrews (Mammalia, Macroscelididae), and a new host and locality record for *Macroscelidopsylla albertyni* De Meillon & Marcus, 1958 (Siphonaptera, Chimaeropsyllidae)—J-C. BEAUCOURNU, I.G. HORAK and L.J. FOURIE 251–253

HUCHZERMEYER, K.D.A.

Clinical and pathological observations on *Streptococcus* sp. infection on South African trout farms with gas supersaturated water supplies—K.D.A. HUCHZERMEYER 95–105

INOKUMA, H.

First evidence of *Anaplasma platys* in *Rhipicephalus sanguineus* (Acari: Ixodida) collected from dogs in Africa—Y.O. SANOGO, B. DAVOUST, H. INOKUMA, J-L. CAMICAS, P. PAROLA and P. BROUQUI 205–212

JONSSON, N.N.

A comparative study on the clinical, parasitological and molecular diagnosis of bovine trypanosomosis in Uganda—J.W. MAGONA, J.S.P. MAYENDE, W. OLAHO-MUKANI, P.G. COLEMAN, N.N. JONSSON, S.C. WELBURN and M.C. EISLER 213–218

JOSEMANS, A.I.

In vitro infection by *Ehrlichia ruminantium* of baby hamster kidney (BHK), Chinese hamster ovary (CHO-K1) and Madin Darby bovine kidney (MDBK) cells—E. ZWEYGARTH and A.I. JOSEMANS 165–168

JOUBERT, J.P.J.

Conditioned feed aversion as a means to prevent tulp (*Homeria pallida*) poisoning in cattle—L.D. SNYMAN, R.A. SCHULTZ, J.P.J. JOUBERT, K.M. BASSON and L. LABUSCHAGNE 43–48

KABONGO, N.

Molecular analysis of bovine viral diarrhoea virus isolates from South Africa—N. KABONGO, C. BAULE and M. VAN VUUREN 273–279

KANIA, S.A.

Evaluation of an indirect enzyme-linked immunosorbent assay for the detection of feline lentivirus-reactive antibodies in wild felids, employing a puma lentivirus-derived synthetic peptide antigen—M. VAN VUUREN, E. STYLIANIDES (nèe DE KLERK), S.A. KANIA, E.E. ZUCKERMAN and W.D. HARDY Jr 1–6

KATJIVENA, H.

Seasonal evolution of faecal egg output by gastrointestinal worms in goats on communal farms in eastern Namibia—F.F. KUMBA, H. KATJIVENA, G. KAUTA and E. LUTAAYA 265–271

KAUTA, G.

Seasonal evolution of faecal egg output by gastrointestinal worms in goats on communal farms in eastern Namibia—F.F. KUMBA, H. KATJIVENA, G. KAUTA and E. LUTAAYA 265–271

KER, J.

Ventriculo-atrial conduction in the ovine heart, caused by premature ventricular complexes—J. KER and E.C. WEBB 107–111

The heart remembers: observations of cardiac memory in the Dorper sheep heart—J. KER, E.C. WEBB, J.A. KER and P.A. BEKKER 299–305

KER, J.A.

The heart remembers: observations of cardiac memory in the Dorper sheep heart—J. KER, E.C. WEBB, J.A. KER and P.A. BEKKER 299–305

KNOBEL, D.L.

Seroconversion in captive African wild dogs (*Lycaon pictus*) following administration of a chicken head bait/SAG-2 oral rabies vaccine combination—D.L. KNOBEL, A. LIEBENBERG and J.T. DU TOIT 73–77

KUMBA, F.F.

Seasonal evolution of faecal egg output by gastrointestinal worms in goats on communal farms in eastern Namibia—F.F. KUMBA, H. KATJIVENA, G. KAUTA and E. LUTAAYA 265–271

LABUSCHAGNE, L.

Conditioned feed aversion as a means to prevent tulp (*Homeria pallida*) poisoning in cattle—L.D. SNYMAN, R.A. SCHULTZ, J.P.J. JOUBERT, K.M. BASSON and L. LABUSCHAGNE 43–48

LIEBENBERG, A.

Seroconversion in captive African wild dogs (*Lycaon pictus*) following administration of a chicken head bait/SAG-2 oral rabies vaccine combination—D.L. KNOBEL, A. LIEBENBERG and J.T. DU TOIT 73–77

LUTAAYA, E.

Seasonal evolution of faecal egg output by gastrointestinal worms in goats on communal farms in eastern Namibia—F.F. KUMBA, H. KATJIVENA, G. KAUTA and E. LUTAAYA 265–271

MACIVOR, K.M. DE F.

Ixodid ticks of Angora and Boer goats, grysbok, common duikers, kudus and scrub hares in Valley Bushveld in the Eastern Cape Province, South Africa—K.M. DE F. MACIVOR and I.G. HORAK 113–120

MAGONA, J.W.

A comparative study on the clinical, parasitological and molecular diagnosis of bovine trypanosomosis in Uganda—J.W. MAGONA, J.S.P. MAYENDE, W. OLAHO-MUKANI, P.G. COLEMAN, N.N. JONSSON, S.C. WELBURN and M.C. EISLER 213–218

MAHAN, S.M.

Comparing the detection of exposure to *Ehrlichia ruminantium* infection on a heartwater-endemic farm by the pCS20 polymerase chain reaction assay and an indirect MAP1-B enzyme linked immunosorbent assay—B.H. SIMBI, T.F. PETER, M.J. BURRIDGE and S.M. MAHAN 231–235

MAROBELA-RABOROKGWE, C.

Comparison of complement fixation test, immunoblotting, indirect ELISA, and competitive ELISA for detecting antibodies to *Mycoplasma mycoides* subspecies *mycoides* small colony (SC) in naturally infected cattle from the 1995 outbreak in Botswana—C. MAROBELA-RABOROKGWE, R. NICHOLAS, R. AYLING and J.B. BASHIRUDDIN . 21–27

MATENGA, ELIZABETH

Effect of host age in the distribution of adult *Trichinella zimbabwensis* in the small intestines of golden hamsters (*Mesocricetus auratus*) and Balb C mice—S. MUKARATIRWA, E. NKULUNGO, ELIZABETH MATENGA and E. BHEBHE 169–173

MATETE, G.

Occurrence, clinical manifestation and the epidemiological implications of naturally occurring canine trypanosomosis in western Kenya—G. MATETE 317–323

MATTHEE, SONJA

Parasites of domestic and wild animals in South Africa. XLIII. Ixodid ticks of domestic dogs and cats in the Western Cape Province—I.G. HORAK and SONJA MATTHEE 187–195

MAYENDE, J.S.P.

A comparative study on the clinical, parasitological and molecular diagnosis of bovine trypanosomosis in Uganda—J.W. MAGONA, J.S.P. MAYENDE, W. OLAHO-MUKANI, P.G. COLEMAN, N.N. JONSSON, S.C. WELBURN and M.C. EISLER 213–218

MBATI, P.A.

Leishmania donovani-derived lipophosphoglycan plus BCG induces a Th1 type immune response but does not protect Syrian golden hamsters (*Mesocricetus auratus*) and BALB/c mice against *Leishmania donovani*—W.K. TONUI, S.S. MPOKE, A.S. ORAGO, S.J. TURCO, P.A. MBATI and G.M. MKOJI 255–263

MCLEOD, A.

Use of rapid rural appraisal and cross-sectional studies in the assessment of constraints in small-holder cattle production systems in the western Kenya highlands—O.S. OKUTHE, A. MCLEOD, J.M. OTTE and G.E. BUYU ... 237–242

MKOJI, G.M.

Leishmania donovani-derived lipophosphoglycan plus BCG induces a Th1 type immune response but does not protect Syrian golden hamsters (*Mesocricetus auratus*) and BALB/c mice against *Leishmania donovani*—W.K. TONUI, S.S. MPOKE, A.S. ORAGO, S.J. TURCO, P.A. MBATI and G.M. MKOJI 255–263

MPOKE, S.S.

Leishmania donovani-derived lipophosphoglycan plus BCG induces a Th1 type immune response but does not protect Syrian golden hamsters (*Mesocricetus auratus*) and BALB/c mice against *Leishmania donovani*—W.K. TONUI, S.S. MPOKE, A.S. ORAGO, S.J. TURCO, P.A. MBATI and G.M. MKOJI 255–263

MUGUNIERI, G.L.

Resistance to trypanocidal drugs—suggestions from field survey on drug use in Kwale district, Kenya—G.L. MUGUNIERI and G.A. MURILLA 29–36

MUKARATIRWA, S.

Effect of host age in the distribution of adult *Trichinella zimbabwensis* in the small intestines of golden hamsters (*Mesocricetus auratus*) and Balb C mice—S. MUKARATIRWA, E. NKULUNGO, ELIZABETH MATENGA and E. BHEBHE 169–173

MURILLA, G.A.

Resistance to trypanocidal drugs—suggestions from field survey on drug use in Kwale district, Kenya—G.L. MUGUNIERI and G.A. MURILLA 29–36

MYBURGH, E.

Review of blackfly (Diptera: Simuliidae) control in South Africa—E. MYBURGH and E.M. NEVILL 307–316

NEL, L.

PCR-based detection of the transovarial transmission of Uruguayan *Babesia bovis* and *Babesia bigemina* vaccine strains—V. GAYO, M. ROMITO, L. NEL, M.A. SOLARI and G.J. VILJOEN 197–204

NEVILL, E.M.

Review of blackfly (Diptera: Simuliidae) control in South Africa—E. MYBURGH and E.M. NEVILL 307–316

NICHOLAS, R.

Comparison of complement fixation test, immunoblotting, indirect ELISA, and competitive ELISA for detecting antibodies to *Mycoplasma mycoides* subspecies *mycoides* small colony (SC) in naturally infected cattle from the 1995 outbreak in Botswana—C. MAROBELA-RABOROKGWE, R. NICHOLAS, R. AYLING and J.B. BASHIRUDDIN . 21–27

NKULUNGO, E.

Effect of host age in the distribution of adult *Trichinella zimbabwensis* in the small intestines of golden hamsters (*Mesocricetus auratus*) and Balb C mice—S. MUKARATIRWA, E. NKULUNGO, ELIZABETH MATENGA and E. BHEBHE 169–173

NYANGE, J.F.C.

Isolation of *Toxoplasma gondii* from goats with a history of reproductive disorders and the prevalence of *Toxoplasma* and chlamydial antibodies—S.P. SHARMA, E.K. BAIPOLEDI, J.F.C. NYANGE and L. TLAGAE 65–68

OHALE, L.O.C.

The morphological characteristics of the antebrachiocarpal joint of the cheetah (*Acinonyx jubatus*)—L.O.C. OHALE and H.B. GROENEWALD 15–20

OKUTHE, O.S.

Use of rapid rural appraisal and cross-sectional studies in the assessment of constraints in small-holder cattle production systems in the western Kenya highlands—O.S. OKUTHE, A. MCLEOD, J.M. OTTE and G.E. BUYU ... 237–242

OLAHO-MUKANI, W.

A comparative study on the clinical, parasitological and molecular diagnosis of bovine trypanosomosis in Uganda—J.W. MAGONA, J.S.P. MAYENDE, W. OLAHO-MUKANI, P.G. COLEMAN, N.N. JONSSON, S.C. WELBURN and M.C. EISLER 213–218

ORAGO, A.S.

Leishmania donovani-derived lipophosphoglycan plus BCG induces a Th1 type immune response but does not protect Syrian golden hamsters (*Mesocricetus auratus*) and BALB/c mice against *Leishmania donovani*—W.K. TONUI, S.S. MPOKE, A.S. ORAGO, S.J. TURCO, P.A. MBATI and G.M. MKOJI 255–263

OTTE, J.M.

Use of rapid rural appraisal and cross-sectional studies in the assessment of constraints in small-holder cattle production systems in the western Kenya highlands—O.S. OKUTHE, A. MCLEOD, J.M. OTTE and G.E. BUYU ... 237–242

PAROLA, P.

First evidence of *Anaplasma platys* in *Rhipicephalus sanguineus* (Acari: Ixodida) collected from dogs in Africa—Y.O. SANOGO, B. DAVOUST, H. INOKUMA, J-L. CAMICAS, P. PAROLA and P. BROUQUI 205–212

PAWESKA, J.T.

Indirect enzyme-linked immunosorbent assay for the detection of antibody against Rift Valley fever virus in domestic and wild ruminant sera—J.T. PAWESKA, S.J. SMITH, I.M. WRIGHT, R. WILLIAMS, A.S. COHEN, A.A. VAN DIJK, A.A. GROBBELAAR, J.E. CROFT, R. SWANEPOEL and G.H. GERDES 49–64

PETER, T.F.

Comparing the detection of exposure to *Ehrlichia ruminantium* infection on a heartwater-endemic farm by the pCS20 polymerase chain reaction assay and an indirect MAP1-B enzyme linked immunosorbent assay—B.H. SIMBI, T.F. PETER, M.J. BURRIDGE and S.M. MAHAN 231–235

PLUMSTEAD, P.

Continuous disinfection as a means to control infectious diseases in poultry. Evaluation of a continuous disinfection programme for broilers–R.R. BRAGG and P. PLUMSTEAD 219–229

PUTTERILL, J.F.

Studies on the genus *Setaria* Viborg, 1795 in South Africa. II. *Setaria scalprum* (Von Linstow, 1908) and *Setaria saegeri* (Le Van Hoa, 1961)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 7–13

General morphology of the oral cavity of the Nile crocodile, *Crocodylus niloticus* (Laurenti, 1768). I. Palate and gingivae—J.F. PUTTERILL and J.T. SOLEY 281–297

ROMITO, M.

PCR-based detection of the transovarial transmission of Uruguayan *Babesia bovis* and *Babesia bigemina* vaccine strains—V. GAYO, M. ROMITO, L. NEL, M.A. SOLARI and G.J. VILJOEN 197–204

SANOGO, Y.O.

First evidence of *Anaplasma platys* in *Rhipicephalus sanguineus* (Acari: Ixodida) collected from dogs in Africa—Y.O. SANOGO, B. DAVOUST, H. INOKUMA, J-L. CAMICAS, P. PAROLA and P. BROUQUI 205–212

SCHULTZ, R.A.

Conditioned feed aversion as a means to prevent tulp (*Homeria pallida*) poisoning in cattle—L.D. SNYMAN, R.A. SCHULTZ, J.P.J. JOUBERT, K.M. BASSON and L. LABUSCHAGNE 43–48

SHARMA, S.P.

Isolation of *Toxoplasma gondii* from goats with a history of reproductive disorders and the prevalence of *Toxoplasma* and chlamydial antibodies—S.P. SHARMA, E.K. BAIPOLEDI, J.F.C. NYANGE and L. TLAGAE 65–68

SIMBI, B.H.

Comparing the detection of exposure to *Ehrlichia ruminantium* infection on a heartwater-endemic farm by the pCS20 polymerase chain reaction assay and an indirect MAP1-B enzyme linked immunosorbent assay—B.H. SIMBI, T.F. PETER, M.J. BURRIDGE and S.M. MAHAN 231–235

SMITH, S.J.

Indirect enzyme-linked immunosorbent assay for the detection of antibody against Rift Valley fever virus in domestic and wild ruminant sera—J.T. PAWESKA, S.J. SMITH, I.M. WRIGHT, R. WILLIAMS, A.S. COHEN, A.A. VAN DIJK, A.A. GROBBELAAR, J.E. CROFT, R. SWANEPOEL and G.H. GERDES 49–64

SNYMAN, L.D.

Conditioned feed aversion as a means to prevent tulp (*Homeria pallida*) poisoning in cattle—L.D. SNYMAN, R.A. SCHULTZ, J.P.J. JOUBERT, K.M. BASSON and L. LABUSCHAGNE 43–48

SOLARI, M.A.

PCR-based detection of the transovarial transmission of Uruguayan *Babesia bovis* and *Babesia bigemina* vaccine strains—V. GAYO, M. ROMITO, L. NEL, M.A. SOLARI and G.J. VILJOEN 197–204

SOLEY, J.T.

General morphology of the oral cavity of the Nile crocodile, *Crocodylus niloticus* (Laurenti, 1768). I. Palate and gingivae—J.F. PUTTERILL and J.T. SOLEY 281–297

STYLIANIDES, E.

Evaluation of an indirect enzyme-linked immunosorbent assay for the detection of feline lentivirus-reactive antibodies in wild felids, employing a puma lentivirus-derived synthetic peptide antigen—M. VAN VUUREN, E. STYLIANIDES (née DE KLERK), S.A. KANIA, E.E. ZUCKERMAN and W.D. HARDY Jr 1–6

SWANEPOEL, R.

Indirect enzyme-linked immunosorbent assay for the detection of antibody against Rift Valley fever virus in domestic and wild ruminant sera—J.T. PAWESKA, S.J. SMITH, I.M. WRIGHT, R. WILLIAMS, A.S. COHEN, A.A. VAN DIJK, A.A. GROBBELAAR, J.E. CROFT, R. SWANEPOEL and G.H. GERDES 49–64

TLAGAE, L.

Isolation of *Toxoplasma gondii* from goats with a history of reproductive disorders and the prevalence of *Toxoplasma* and chlamydial antibodies—S.P. SHARMA, E.K. BAIPOLEDI, J.F.C. NYANGE and L. TLAGAE 65–68

TONUI, W.K.

Leishmania donovani-derived lipophosphoglycan plus BCG induces a Th1 type immune response but does not protect Syrian golden hamsters (*Mesocricetus auratus*) and BALB/c mice against *Leishmania donovani*—W.K. TONUI, S.S. MPOKE, A.S. ORAGO, S.J. TURCO, P.A. MBATI and G.M. MKOJI 255–263

TURCO, S.J.

Leishmania donovani-derived lipophosphoglycan plus BCG induces a Th1 type immune response but does not protect Syrian golden hamsters (*Mesocricetus auratus*) and BALB/c mice against *Leishmania donovani*—W.K. TONUI, S.S. MPOKE, A.S. ORAGO, S.J. TURCO, P.A. MBATI and G.M. MKOJI 255–263

VAN DIJK, A.A.

Indirect enzyme-linked immunosorbent assay for the detection of antibody against Rift Valley fever virus in domestic and wild ruminant sera—J.T. PAWESKA, S.J. SMITH, I.M. WRIGHT, R. WILLIAMS, A.S. COHEN, A.A. VAN DIJK, A.A. GROBBELAAR, J.E. CROFT, R. SWANEPOEL and G.H. GERDES 49–64

VAN NIEKERK, D.J.

Population structure of *Argas arboreus* (Acari: Argasidae) ticks associated with seasonally abandoned mixed heronries, dominated by cattle egrets (*Bubulcus ibis*), in South Africa—V.N. BELOZEROV, D.J. VAN NIEKERK and H.J.B. BUTLER 325–330

VAN RENSBURG, L.J.

Schistosoma mattheei—an ovum containing twin miracidia—L.J. VAN RENSBURG and J.A. VAN WYK 69–71

Description and classification of abnormal cercariae of *Schistosoma mattheei* and a method of concentration and collection—J.A. VAN WYK and L.J. VAN RENSBURG 121–129

VAN VUUREN, M.

Evaluation of an indirect enzyme-linked immunosorbent assay for the detection of feline lentivirus-reactive antibodies in wild felids, employing a puma lentivirus-derived synthetic peptide antigen—M. VAN VUUREN, E. STYLIANIDES (née DE KLERK), S.A. KANIA, E.E. ZUCKERMAN and W.D. HARDY Jr 1–6

Molecular analysis of bovine viral diarrhoea virus isolates from South Africa—N. KABONGO, C. BAULE and M. VAN VUUREN 273–279

VAN WYK, J.A.

Schistosoma mattheei—an ovum containing twin miracidia—L.J. VAN RENSBURG and J.A. VAN WYK 69–71

Description and classification of abnormal cercariae of *Schistosoma mattheei* and a method of concentration and collection—J.A. VAN WYK and L.J. VAN RENSBURG 121–129

VILJOEN, G.J.

PCR-based detection of the transovarial transmission of Uruguayan *Babesia bovis* and *Babesia bigemina* vaccine strains—V. GAYO, M. ROMITO, L. NEL, M.A. SOLARI and G.J. VILJOEN 197–204

WATERMEYER, R.

Studies on the genus *Setaria* Viborg, 1795 in South Africa. II. *Setaria scalprum* (Von Linstow, 1908) and *Setaria saegeri* (Le Van Hoa, 1961)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 7–13

WEBB, E.C.

Ventriculo-atrial conduction in the ovine heart, caused by premature ventricular complexes—J. KER and E.C. WEBB 107–111

The heart remembers: observations of cardiac memory in the Dorper sheep heart—J. KER, E.C. WEBB, J.A. KER and P.A. BEKKER 299–305

WELBURN, S.C.

A comparative study on the clinical, parasitological and molecular diagnosis of bovine trypanosomosis in Uganda—J.W. MAGONA, J.S.P. MAYENDE, W. OLAHO-MUKANI, P.G. COLEMAN, N.N. JONSSON, S.C. WELBURN and M.C. EISLER 213–218

WILLIAMS, R.

Indirect enzyme-linked immunosorbent assay for the detection of antibody against Rift Valley fever virus in domestic and wild ruminant sera—J.T. PAWESKA, S.J. SMITH, I.M. WRIGHT, R. WILLIAMS, A.S. COHEN, A.A. VAN DIJK, A.A. GROBBELAAR, J.E. CROFT, R. SWANEPOEL and G.H. GERDES 49–64

WRIGHT, I.M.

Indirect enzyme-linked immunosorbent assay for the detection of antibody against Rift Valley fever virus in domestic and wild ruminant sera—J.T. PAWESKA, S.J. SMITH, I.M. WRIGHT, R. WILLIAMS, A.S. COHEN, A.A. VAN DIJK, A.A. GROBBELAAR, J.E. CROFT, R. SWANEPOEL and G.H. GERDES 49–64

ZUCKERMAN, E.E.

Evaluation of an indirect enzyme-linked immunosorbent assay for the detection of feline lentivirus-reactive antibodies in wild felids, employing a puma lentivirus-derived synthetic peptide antigen—M. VAN VUUREN, E. STYLIANIDES (née DE KLERK), S.A. KANIA, E.E. ZUCKERMAN and W.D. HARDY Jr 1–6

ZWEYGARTH, E.

In vitro infection by *Ehrlichia ruminantium* of baby hamster kidney (BHK), Chinese hamster ovary (CHO-K1) and Madin Darby bovine kidney (MDBK) cells—E. ZWEYGARTH and A.I. JOSEMANS 165–168

ONDERSTEPOORT JOURNAL OF VETERINARY RESEARCH

Volume 70 • Numbers 1–4 • 2003

Subject index

Acari: Argasidae

Population structure of *Argas arboreus* (Acari: Argasidae) ticks associated with seasonally abandoned mixed heronries, dominated by cattle egrets (*Bubulcus ibis*), in South Africa—V.N. BELOZEROV, D.J. VAN NIEKERK and H.J.B. BUTLER 325–330

Acari: Ixodidae

Climate change decreases habitat suitability for some tick species (Acari: Ixodidae) in South Africa—A. ESTRADA-PEÑA 79–93

First evidence of *Anaplasma platys* in *Rhipicephalus sanguineus* (Acari: Ixodida) collected from dogs in Africa—Y.O. SANOGO, B. DAVOUST, H. INOKUMA, J-L. CAMICAS, P. PAROLA and P. BROUQUI 205–212

Acinonyx jubatus

The morphological characteristics of the antebrachiocarpal joint of the cheetah (*Acinonyx jubatus*)—L.O.C. OHALE and H.B. GROENEWALD 15–20

Aepyceros melampus

Parasites of domestic and wild animals in South Africa. XLI. Arthropod parasites of impalas, *Aepyceros melampus* in the Kruger National Park—I.G. HORAK, G.J. GALLIVAN, L.E.O. BRAACK, J. BOOMKER and V. DE VOS 131–163

Amblyomma hebraeum

Comparing the detection of exposure to *Ehrlichia ruminantium* infection on a heartwater-endemic farm by the pCS20 polymerase chain reaction assay and an indirect MAP1-B enzyme linked immunosorbent assay—B.H. SIMBI, T.F. PETER, M.J. BURRIDGE and S.M. MAHAN 231–235

Anaplasma platys

First evidence of *Anaplasma platys* in *Rhipicephalus sanguineus* (Acari: Ixodida) collected from dogs in Africa—Y.O. SANOGO, B. DAVOUST, H. INOKUMA, J-L. CAMICAS, P. PAROLA and P. BROUQUI 205–212

Argas arboreus

Population structure of *Argas arboreus* (Acari: Argasidae) ticks associated with seasonally abandoned mixed heronries, dominated by cattle egrets (*Bubulcus ibis*), in South Africa—V.N. BELOZEROV, D.J. VAN NIEKERK and H.J.B. BUTLER 325–330

Arthropod parasites

Parasites of domestic and wild animals in South Africa. XLI. Arthropod parasites of impalas, *Aepyceros melampus* in the Kruger National Park—I.G. HORAK, G.J. GALLIVAN, L.E.O. BRAACK, J. BOOMKER and V. DE VOS 131–163

Babesia bigemina

PCR-based detection of the transovarial transmission of Uruguayan *Babesia bovis* and *Babesia bigemina* vaccine strains—V. GAYO, M. ROMITO, L. NEL, M.A. SOLARI and G.J. VILJOEN 197–204

Babesia bovis

PCR-based detection of the transovarial transmission of Uruguayan *Babesia bovis* and *Babesia bigemina* vaccine strains—V. GAYO, M. ROMITO, L. NEL, M.A. SOLARI and G.J. VILJOEN 197–204

Bacille Calmette-Guérin (BCG)

Leishmania donovani-derived lipophosphoglycan plus BCG induces a Th1 type immune response but does not protect Syrian golden hamsters (*Mesocricetus auratus*) and BALB/c mice against *Leishmania donovani*—W.K. TONUI, S.S. MPOKE, A.S. ORAGO, S.J. TURCO, P.A. MBATI and G.M. MKOJI 255–263

Bacillus thuringiensis* var. *israelensis

Review of blackfly (Diptera: Simuliidae) control in South Africa—E MYBURGH and E.M. NEVILL 307–316

BALB/c

Effect of host age in the distribution of adult *Trichinella zimbabwensis* in the small intestines of golden hamsters (*Mesocricetus auratus*) and Balb C mice—S. MUKARATIRWA, E. NKULUNGO, ELIZABETH MATENGA and E. BHEBHE 169–173

Leishmania donovani-derived lipophosphoglycan plus BCG induces a Th1 type immune response but does not protect Syrian golden hamsters (*Mesocricetus auratus*) and BALB/c mice against *Leishmania donovani*—W.K. TONUI, S.S. MPOKE, A.S. ORAGO, S.J. TURCO, P.A. MBATI and G.M. MKOJI 255–263

Blackfly

Review of blackfly (Diptera: Simuliidae) control in South Africa—E. MYBURGH and E.M. NEVILL 307–316

Boophilus microplus

PCR-based detection of the transovarial transmission of Uruguayan *Babesia bovis* and *Babesia bigemina* vaccine strains—V. GAYO, M. ROMITO, L. NEL, M.A. SOLARI and G.J. VILJOEN 197–204

Botswana

Comparison of complement fixation test, immunoblotting, indirect ELISA, and competitive ELISA for detecting antibodies to *Mycoplasma mycoides* subspecies *mycoides* small colony (SC) in naturally infected cattle from the 1995 outbreak in Botswana—C. MAROBELA-RABOROKGWE, R. NICHOLAS, R. AYLING and J.B. BASHIRUDDIN .. 21–27

Bovine pleuropmeumonia

Comparison of complement fixation test, immunoblotting, indirect ELISA, and competitive ELISA for detecting antibodies to *Mycoplasma mycoides* subspecies *mycoides* small colony (SC) in naturally infected cattle from the 1995 outbreak in Botswana—C. MAROBELA-RABOROKGWE, R. NICHOLAS, R. AYLING and J.B. BASHIRUDDIN .. 21–27

Bovine trypanosomosis

A comparative study on the clinical, parasitological and molecular diagnosis of bovine trypanosomosis in Uganda—J.W. MAGONA, J.S.P. MAYENDE, W. OLAHO-MUKANI, P.G. COLEMAN, N.N. JONSSON, S.C. WELBURN and M.C. EISLER 213–218

Bovine viral diarrhoea virus

Molecular analysis of bovine viral diarrhoea virus isolates from South Africa—N. KABONGO, C. BAULE and M. VAN VUUREN 273–279

Broilers

Continuous disinfection as a means to control infectious diseases in poultry. Evaluation of a continuous disinfection programme for broilers–R.R. BRAGG and P. PLUMSTEAD 219–229

Bubulcus ibis

Population structure of *Argas arboreus* (Acari: Argasidae) ticks associated with seasonally abandoned mixed heronries, dominated by cattle egrets (*Bubulcus ibis*), in South Africa—V.N. BELOZEROV, D.J. VAN NIEKERK and H.J.B. BUTLER 325–330

Canine trypanosomosis

Occurrence, clinical manifestation and the epidemiological implications of naturally occurring canine trypanosomosis in western Kenya—G. MATETE 317–323

Cardiac memory

The heart remembers: observations of cardiac memory in the Dorper sheep heart—J. KER, E.C. WEBB, J.A. KER and P.A. BEKKER 299–305

Cats

Parasites of domestic and wild animals in South Africa. XLIII. Ixodid ticks of domestic dogs and cats in the Western Cape Province—I.G. HORAK and SONJA MATTHEE 187–195

Cattle

Comparison of complement fixation test, immunoblotting, indirect ELISA, and competitive ELISA for detecting antibodies to *Mycoplasma mycoides* subspecies *mycoides* small colony (SC) in naturally infected cattle from the 1995 outbreak in Botswana—C. MAROBELA-RABOROKGWE, R. NICHOLAS, R. AYLING and J.B. BASHIRUDDIN .. 21–27

Conditioned feed aversion as a means to prevent tulp (*Homeria pallida*) poisoning in cattle—L.D. SNYMAN, R.A. SCHULTZ, J.P.J. JOUBERT, K.M. BASSON and L. LABUSCHAGNE 43–48

A comparative study on the clinical, parasitological and molecular diagnosis of bovine trypanosomosis in Uganda—J.W. MAGONA, J.S.P. MAYENDE, W. OLAHO-MUKANI, P.G. COLEMAN, N.N. JONSSON, S.C. WELBURN and M.C. EISLER 213–218

Use of rapid rural appraisal and cross-sectional studies in the assessment of constraints in small-holder cattle production systems in the western Kenya highlands—O.S. OKUTHE, A. MCLEOD, J.M. OTTE and G.E. BUYU 237–242

Molecular analysis of bovine viral diarrhoea virus isolates from South Africa—N. KABONGO, C. BAULE and M. VAN VUUREN 273–279

Cercariae

Description and classification of abnormal cercariae of *Schistosoma mattheei* and a method of concentration and collection—J.A. VAN WYK and L.J. VAN RENSBURG 121–129

Cheetah

The morphological characteristics of the antebrachiocarpal joint of the cheetah (*Acinonyx jubatus*)—L.O.C. OHALE and H.B. GROENEWALD 15–20

Chlamydia psittaci

Isolation of *Toxoplasma gondii* from goats with a history of reproductive disorders and the prevalence of *Toxoplasma* and chlamydial antibodies—S.P. SHARMA, E.K. BAIPOLEDI, J.F.C. NYANGE and L. TLAGAE 65–68

Complement fixation test

Comparison of complement fixation test, immunoblotting, indirect ELISA, and competitive ELISA for detecting antibodies to *Mycoplasma mycoides* subspecies *mycoides* small colony (SC) in naturally infected cattle from the 1995 outbreak in Botswana—C. MAROBELA-RABOROKGWE, R. NICHOLAS, R. AYLING and J.B. BASHIRUDDIN .. 21–27

***Crocodylus niloticus* (Laurenti, 1768)**

General morphology of the oral cavity of the Nile crocodile, *Crocodylus niloticus* (Laurenti, 1768). I. Palate and gingivae—J.F. PUTTERILL and J.T. SOLEY 281–297

Demeillonia granti

Fleas of elephant shrews (Mammalia, Macroscelididae), and a new host and locality record for *Macroscelidopsylla albertyni* De Meillon & Marcus, 1958 (Siphonaptera, Chimaeropsyllidae)—J-C. BEAUCOURNU, I.G. HORAK and L.J. FOURIE 251–253

Democratic Republic of Congo

First evidence of *Anaplasma platys* in *Rhipicephalus sanguineus* (Acari: Ixodida) collected from dogs in Africa—Y.O. SANOGO, B. DAVOUST, H. INOKUMA, J-L. CAMICAS, P. PAROLA and P. BROUQUI 205–212

Diptera: Simuliidae

Review of blackfly (Diptera: Simuliidae) control in South Africa—E. MYBURGH and E.M. NEVILL 307–316

Dogs

Seroconversion in captive African wild dogs (*Lycaon pictus*) following administration of a chicken head bait/SAG-2 oral rabies vaccine combination—D.L. KNOBEL, A. LIEBENBERG and J.T. DU TOIT 73–77

Parasites of domestic and wild animals in South Africa. XLIII. Ixodid ticks of domestic dogs and cats in the Western Cape Province—I.G. HORAK and SONJA MATTHEE 187–195

First evidence of *Anaplasma platys* in *Rhipicephalus sanguineus* (Acari: Ixodida) collected from dogs in Africa—Y.O. SANOGO, B. DAVOUST, H. INOKUMA, J-L. CAMICAS, P. PAROLA and P. BROUQUI 205–212

Duikers

Ixodid ticks of Angora and Boer goats, grysbok, common duikers, kudus and scrub hares in Valley Bushveld in the Eastern Cape Province, South Africa—K.M. DE F. MACIVOR and I.G. HORAK 113–120

Egrets

Population structure of *Argas arboreus* (Acari: Argasidae) ticks associated with seasonally abandoned mixed heronries, dominated by cattle egrets (*Bubulcus ibis*), in South Africa—V.N. BELOZEROV, D.J. VAN NIEKERK and H.J.B. BUTLER 325–330

Ehrlichia ruminantium

In vitro infection by *Ehrlichia ruminantium* of baby hamster kidney (BHK), Chinese hamster ovary (CHO-K1) and Madin Darby bovine kidney (MDBK) cells—E. ZWEYGARTH and A.I. JOSEMANS 165–168

Comparing the detection of exposure to *Ehrlichia ruminantium* infection on a heartwater-endemic farm by the pCS20 polymerase chain reaction assay and an indirect MAP1-B enzyme linked immunosorbent assay—B.H. SIMBI, T.F. PETER, M.J. BURRIDGE and S.M. MAHAN 231–235

Detection of antibodies to the *Ehrlichia ruminantium* MAP1-B antigen in goat sera from three communal land areas of Zimbabwe by an indirect enzyme-linked immunosorbent assay—O. KAKONO, T. HOVE, D. GEYSEN and S. MAHAN 243–249

Elephant shrew

Fleas of elephant shrews (Mammalia, Macroscelididae), and a new host and locality record for *Macroscelidopsylla albertyni* De Meillon & Marcus, 1958 (Siphonaptera, Chimaeropsyllidae)—J-C. BEAUCOURNU, I.G. HORAK and L.J. FOURIE 251–253

Elephantulus myurus* and *Elephantulus edwardii

Fleas of elephant shrews (Mammalia, Macroscelididae), and a new host and locality record for *Macroscelidopsylla albertyni* De Meillon & Marcus, 1958 (Siphonaptera, Chimaeropsyllidae)—J-C. BEAUCOURNU, I.G. HORAK and L.J. FOURIE 251–253

Enzyme-linked immunosorbent assay (ELISA)

Evaluation of an indirect enzyme-linked immunosorbent assay for the detection of feline lentivirus-reactive antibodies in wild felids, employing a puma lentivirus-derived synthetic peptide antigen—M. VAN VUUREN, E. STYLIANIDES (nèe DE KLERK), S.A. KANIA, E.E. ZUCKERMAN and W.D. HARDY Jr 1–6

Comparison of complement fixation test, immunoblotting, indirect ELISA, and competitive ELISA for detecting antibodies to *Mycoplasma mycoides* subspecies *mycoides* small colony (SC) in naturally infected cattle from the 1995 outbreak in Botswana—C. MAROBELA-RABOROKGWE, R. NICHOLAS, R. AYLING and J.B. BASHIRUDDIN .. 21–27

Indirect enzyme-linked immunosorbent assay for the detection of antibody against Rift Valley fever virus in domestic and wild ruminant sera—J.T. PAWESKA, S.J. SMITH, I.M. WRIGHT, R. WILLIAMS, A.S. COHEN, A.A. VAN DIJK, A.A. GROBBELAAR, J.E. CROFT, R. SWANEPOEL and G.H. GERDES 49–64

Comparing the detection of exposure to *Ehrlichia ruminantium* infection on a heartwater-endemic farm by the pCS20 polymerase chain reaction assay and an indirect MAP1-B enzyme linked immunosorbent assay—B.H. SIMBI, T.F. PETER, M.J. BURRIDGE and S.M. MAHAN 231–235

Detection of antibodies to the *Ehrlichia ruminantium* MAP1-B antigen in goat sera from three communal land areas of Zimbabwe by an indirect enzyme-linked immunosorbent assay—O. KAKONO, T. HOVE, D. GEYSEN and S. MAHAN 243–249

Faecal egg output

Seasonal evolution of faecal egg output by gastrointestinal worms in goats on communal farms in eastern Namibia—F.F. KUMBA, H. KATJIVENA, G. KAUTA and E. LUTAAYA 265–271

FAMACHA© system

Seasonal evolution of faecal egg output by gastrointestinal worms in goats on communal farms in eastern Namibia—F.F. KUMBA, H. KATJIVENA, G. KAUTA and E. LUTAAYA 265–271

Feline immunodeficiency virus (FIV)

Evaluation of an indirect enzyme-linked immunosorbent assay for the detection of feline lentivirus-reactive antibodies in wild felids, employing a puma lentivirus-derived synthetic peptide antigen—M. VAN VUUREN, E. STYLIANIDES (nèe DE KLERK), S.A. KANIA, E.E. ZUCKERMAN and W.D. HARDY Jr 1–6

Filarids

Studies on the genus *Setaria* Viborg, 1795 in South Africa. II. *Setaria scalprum* (Von Linstow, 1908) and *Setaria saegeri* (Le Van Hoa, 1961)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 7–13

Fleas

Fleas of elephant shrews (Mammalia, Macroscelididae), and a new host and locality record for *Macroscelidopsylla albertyni* De Meillon & Marcus, 1958 (Siphonaptera, Chimaeropsyllidae)—J-C. BEAUCOURNU, I.G. HORAK and L.J. FOURIE 251–253

Gas bubble disease

Clinical and pathological observations on *Streptococcus* sp. infection on South African trout farms with gas supersaturated water supplies—K.D.A. HUCHZERMEYER 95–105

Gastrointestinal nematodes/worms

Seasonal evolution of faecal egg output by gastrointestinal worms in goats on communal farms in eastern Namibia—F.F. KUMBA, H. KATJIVENA, G. KAUTA and E. LUTAAYA 265–271

Goats

Isolation of *Toxoplasma gondii* from goats with a history of reproductive disorders and the prevalence of Toxoplasma and chlamydial antibodies—S.P. SHARMA, E.K. BAIPOLEDI, J.F.C. NYANGE and L. TLAGAE 65–68

Ixodid ticks of Angora and Boer goats, grysbok, common duikers, kudus and scrub hares in Valley Bushveld in the Eastern Cape Province, South Africa—K.M. DE F. MACIVOR and I.G. HORAK 113–120

Detection of antibodies to the *Ehrlichia ruminantium* MAP1-B antigen in goat sera from three communal land areas of Zimbabwe by an indirect enzyme-linked immunosorbent assay—O. KAKONO, T. HOVE, D. GEYSEN and S. MAHAN 243–249

Seasonal evolution of faecal egg output by gastrointestinal worms in goats on communal farms in eastern Namibia—F.F. KUMBA, H. KATJIVENA, G. KAUTA and E. LUTAAYA 265–271

Grey rhebuck

Parasites of South African wildlife. XVII. *Ostertagia triquetra* n. sp. (Nematoda: Trichostrongylina) from the grey rhebuck, *Pelea capreolus* (Forster, 1790)—J. BOOMKER and M-C. DURETTE-DESSET 37–41

Grysbok

Ixodid ticks of Angora and Boer goats, grysbok, common duikers, kudus and scrub hares in Valley Bushveld in the Eastern Cape Province, South Africa—K.M. DE F. MACIVOR and I.G. HORAK 113–120

Haemaphysalis leachi

Parasites of domestic and wild animals in South Africa. XLIII. Ixodid ticks of domestic dogs and cats in the Western Cape Province—I.G. HORAK and SONJA MATTHEE 187–195

Hamsters

In vitro infection by *Ehrlichia ruminantium* of baby hamster kidney (BHK), Chinese hamster ovary (CHO-K1) and Madin Darby bovine kidney (MDBK) cells—E. ZWEYGARTH and A.I. JOSEMANS 165–168

Effect of host age in the distribution of adult *Trichinella zimbabwensis* in the small intestines of golden hamsters (*Mesocricetus auratus*) and Balb C mice—S. MUKARATIRWA, E. NKULUNGO, ELIZABETH MATENGA and E. BHEBHE 169–173

Leishmania donovani-derived lipophosphoglycan plus BCG induces a Th1 type immune response but does not protect Syrian golden hamsters (*Mesocricetus auratus*) and BALB/c mice against *Leishmania donovani*—W.K. TONUI, S.S. MPOKE, A.S. ORAGO, S.J. TURCO, P.A. MBATI and G.M. MKOJI 255–263

Heart

Ventriculo-atrial conduction in the ovine heart, caused by premature ventricular complexes—J. KER and E.C. WEBB 107–111

The heart remembers: observations of cardiac memory in the Dorper sheep heart—J. KER, E.C. WEBB, J.A. KER and P.A. BEKKER 299–305

Heartwater

In vitro infection by *Ehrlichia ruminantium* of baby hamster kidney (BHK), Chinese hamster ovary (CHO-K1) and Madin Darby bovine kidney (MDBK) cells—E. ZWEYGARTH and A.I. JOSEMANS 165–168

Comparing the detection of exposure to *Ehrlichia ruminantium* infection on a heartwater-endemic farm by the pCS20 polymerase chain reaction assay and an indirect MAP1-B enzyme linked immunosorbent assay—B.H. SIMBI, T.F. PETER, M.J. BURRIDGE and S.M. MAHAN 231–235

Detection of antibodies to the *Ehrlichia ruminantium* MAP1-B antigen in goat sera from three communal land areas of Zimbabwe by an indirect enzyme-linked immunosorbent assay—O. KAKONO, T. HOVE, D. GEYSEN and S. MAHAN 243–249

Helminths

Parasites of domestic and wild animals in South Africa. XLII. Helminths of sheep on four farms in the Eastern Cape Province—I.G. HORAK 175–186

Heronries

Population structure of *Argas arboreus* (Acari: Argasidae) ticks associated with seasonally abandoned mixed heronries, dominated by cattle egrets (*Bubulcus ibis*), in South Africa—V.N. BELOZEROV, D.J. VAN NIEKERK and H.J.B. BUTLER 325–330

Homeria pallida

Conditioned feed aversion as a means to prevent tulp (*Homeria pallida*) poisoning in cattle—L.D. SNYMAN, R.A. SCHULTZ, J.P.J. JOUBERT, K.M. BASSON and L. LABUSCHAGNE 43–48

Immunoblotting

Comparison of complement fixation test, immunoblotting, indirect ELISA, and competitive ELISA for detecting antibodies to *Mycoplasma mycoides* subspecies *mycoides* small colony (SC) in naturally infected cattle from the 1995 outbreak in Botswana—C. MAROBELA-RABOROKGWE, R. NICHOLAS, R. AYLING and J.B. BASHIRUDDIN .. 21–27

Impalas

Parasites of domestic and wild animals in South Africa. XLI. Arthropod parasites of impalas, *Aepyceros melampus* in the Kruger National Park—I.G. HORAK, G.J. GALLIVAN, L.E.O. BRAACK, J. BOOMKER and V. DE VOS 131–163

Ixodid ticks

Parasites of domestic and wild animals in South Africa. XLIII. Ixodid ticks of domestic dogs and cats in the Western Cape Province—I.G. HORAK and SONJA MATTHEE 187–195

Kenya

Resistance to trypanocidal drugs—suggestions from field survey on drug use in Kwale district, Kenya—G.L. MUGUNIERI and G.A. MURILLA 29–36

Use of rapid rural appraisal and cross-sectional studies in the assessment of constraints in small-holder cattle production systems in the western Kenya highlands—O.S. OKUTHE, A. MCLEOD, J.M. OTTE and G.E. BUYU 237–242

Occurrence, clinical manifestation and the epidemiological implications of naturally occurring canine trypanosomosis in western Kenya—G. MATETE 317–323

Kudus

Ixodid ticks of Angora and Boer goats, grysbok, common duikers, kudus and scrub hares in Valley Bushveld in the Eastern Cape Province, South Africa—K.M. DE F. MACIVOR and I.G. HORAK 113–120

Leishmania donovani

Leishmania donovani-derived lipophosphoglycan plus BCG induces a Th1 type immune response but does not protect Syrian golden hamsters (*Mesocricetus auratus*) and BALB/c mice against *Leishmania donovani*—W.K. TONUI, S.S. MPOKE, A.S. ORAGO, S.J. TURCO, P.A. MBATI and G.M. MKOJI 255–263

Lentivirus

Evaluation of an indirect enzyme-linked immunosorbent assay for the detection of feline lentivirus-reactive antibodies in wild felids, employing a puma lentivirus-derived synthetic peptide antigen—M. VAN VUUREN, E. STYLIANIDES (née DE KLERK), S.A. KANIA, E.E. ZUCKERMAN and W.D. HARDY Jr 1–6

Lice

Parasites of domestic and wild animals in South Africa. XLI. Arthropod parasites of impalas, *Aepyceros melampus* in the Kruger National Park—I.G. HORAK, G.J. GALLIVAN, L.E.O. BRAACK, J. BOOMKER and V. DE VOS 131–163

Lions

Evaluation of an indirect enzyme-linked immunosorbent assay for the detection of feline lentivirus-reactive antibodies in wild felids, employing a puma lentivirus-derived synthetic peptide antigen—M. VAN VUUREN, E. STYLIANIDES (née DE KLERK), S.A. KANIA, E.E. ZUCKERMAN and W.D. HARDY Jr 1–6

Lipophosphoglycan

Leishmania donovani-derived lipophosphoglycan plus BCG induces a Th1 type immune response but does not protect Syrian golden hamsters (*Mesocricetus auratus*) and BALB/c mice against *Leishmania donovani*—W.K. TONUI, S.S. MPOKE, A.S. ORAGO, S.J. TURCO, P.A. MBATI and G.M. MKOJI 255–263

Lycaon pictus

Seroconversion in captive African wild dogs (*Lycaon pictus*) following administration of a chicken head bait/SAG-2 oral rabies vaccine combination—D.L. KNOBEL, A. LIEBENBERG and J.T. DU TOIT 73–77

Macroscelididae

Fleas of elephant shrews (Mammalia, Macroscelididae), and a new host and locality record for *Macroscelidopsylla albertyni* De Meillon & Marcus, 1958 (Siphonaptera, Chimaeropsyllidae)—J-C. BEAUCOURNU, I.G. HORAK and L.J. FOURIE 251–253

Macroscelidopsylla albertyni

Fleas of elephant shrews (Mammalia, Macroscelididae), and a new host and locality record for *Macroscelidopsylla albertyni* De Meillon & Marcus, 1958 (Siphonaptera, Chimaeropsyllidae)—J-C. BEAUCOURNU, I.G. HORAK and L.J. FOURIE 251–253

Madin Darby bovine kidney cells

In vitro infection by *Ehrlichia ruminantium* of baby hamster kidney (BHK), Chinese hamster ovary (CHO-K1) and Madin Darby bovine kidney (MDBK) cells—E. ZWEYGARTH and A.I. JOSEMANS 165–168

MAP1-B

Comparing the detection of exposure to *Ehrlichia ruminantium* infection on a heartwater-endemic farm by the pCS20 polymerase chain reaction assay and an indirect MAP1-B enzyme linked immunosorbent assay—B.H. SIMBI, T.F. PETER, M.J. BURRIDGE and S.M. MAHAN 231–235

Detection of antibodies to the *Ehrlichia ruminantium* MAP1-B antigen in goat sera from three communal land areas of Zimbabwe by an indirect enzyme-linked immunosorbent assay—O. KAKONO, T. HOVE, D. GEYSEN and S. MAHAN 243–249

Mesocricetus auratus

Effect of host age in the distribution of adult *Trichinella zimbabwensis* in the small intestines of golden hamsters (*Mesocricetus auratus*) and Balb C mice—S. MUKARATIRWA, E. NKULUNGO, ELIZABETH MATENGA and E. BHEBHE 169–173

Leishmania donovani-derived lipophosphoglycan plus BCG induces a Th1 type immune response but does not protect Syrian golden hamsters (*Mesocricetus auratus*) and BALB/c mice against *Leishmania donovani*—W.K. TONUI, S.S. MPOKE, A.S. ORAGO, S.J. TURCO, P.A. MBATI and G.M. MKOJI 255–263

Mice

Effect of host age in the distribution of adult *Trichinella zimbabwensis* in the small intestines of golden hamsters (*Mesocricetus auratus*) and Balb C mice—S. MUKARATIRWA, E. NKULUNGO, ELIZABETH MATENGA and E. BHEBHE 169–173

Leishmania donovani-derived lipophosphoglycan plus BCG induces a Th1 type immune response but does not protect Syrian golden hamsters (*Mesocricetus auratus*) and BALB/c mice against *Leishmania donovani*—W.K. TONUI, S.S. MPOKE, A.S. ORAGO, S.J. TURCO, P.A. MBATI and G.M. MKOJI 255–263

Miracidia

Schistosoma mattheei—an ovum containing twin miracidia—L.J. VAN RENSBURG and J.A. VAN WYK 69–71

Mycoplasma mycoides

Comparison of complement fixation test, immunoblotting, indirect ELISA, and competitive ELISA for detecting antibodies to *Mycoplasma mycoides* subspecies *mycoides* small colony (SC) in naturally infected cattle from the 1995 outbreak in Botswana—C. MAROBELA-RABOROKGWE, R. NICHOLAS, R. AYLING and J.B. BASHIRUDDIN .. 21–27

Namibia

Seasonal evolution of faecal egg output by gastrointestinal worms in goats on communal farms in eastern Namibia—F.F. KUMBA, H. KATJIVENA, G. KAUTA and E. LUTAAYA 265–271

Nematoda: Trichostrongylina

Parasites of South African wildlife. XVII. *Ostertagia triquetra* n. sp. (Nematoda: Trichostrongylina) from the grey rhebuck, *Pelea capreolus* (Forster, 1790)—J. BOOMKER and M-C. DURETTE-DESSET 37–41

Newcastle Disease

Continuous disinfection as a means to control infectious diseases in poultry. Evaluation of a continuous disinfection programme for broilers–R.R. BRAGG and P. PLUMSTEAD 219–229

Nile crocodile

General morphology of the oral cavity of the Nile crocodile, *Crocodylus niloticus* (Laurenti, 1768). I. Palate and gingivae—J.F. PUTTERILL and J.T. SOLEY 281–297

Onchorhynchus mykiss

Clinical and pathological observations on *Streptococcus* sp. infection on South African trout farms with gas supersaturated water supplies—K.D.A. HUCHZERMEYER 95–105

Ostertagia triquetra

Parasites of South African wildlife. XVII. *Ostertagia triquetra* n. sp. (Nematoda: Trichostrongylina) from the grey rhebuck, *Pelea capreolus* (Forster, 1790)—J. BOOMKER and M-C. DURETTE-DESSET 37–41

Ovine

Ventriculo-atrial conduction in the ovine heart, caused by premature ventricular complexes—J. KER and E.C. WEBB 107–111

Parasites

Parasites of South African wildlife. XVII. *Ostertagia triquetra* n. sp. (Nematoda: Trichostrongylina) from the grey rhebuck, *Pelea capreolus* (Forster, 1790)—J. BOOMKER and M-C. DURETTE-DESSET 37–41

Parasites of domestic and wild animals in South Africa. XLII. Helminths of sheep on four farms in the Eastern Cape Province—I.G. HORAK 175–186

Parasites of domestic and wild animals in South Africa. XLIII. Ixodid ticks of domestic dogs and cats in the Western Cape Province—I.G. HORAK and SONJA MATTHEE 187–195

pCS20

Comparing the detection of exposure to *Ehrlichia ruminantium* infection on a heartwater-endemic farm by the pCS20 polymerase chain reaction assay and an indirect MAP1-B enzyme linked immunosorbent assay—B.H. SIMBI, T.F. PETER, M.J. BURRIDGE and S.M. MAHAN 231–235

Pelea capreolus

Parasites of South African wildlife. XVII. *Ostertagia triquetra* n. sp. (Nematoda: Trichostrongylina) from the grey rhebuck, *Pelea capreolus* (Forster, 1790)—J. BOOMKER and M-C. DURETTE-DESSET 37–41

Polymerase chain reaction (PCR) assay

PCR-based detection of the transovarial transmission of Uruguayan *Babesia bovis* and *Babesia bigemina* vaccine strains—V. GAYO, M. ROMITO, L. NEL, M.A. SOLARI and G.J. VILJOEN 197–204

Poultry

Continuous disinfection as a means to control infectious diseases in poultry. Evaluation of a continuous disinfection programme for broilers–R.R. BRAGG and P. PLUMSTEAD 219–229

Puma

Evaluation of an indirect enzyme-linked immunosorbent assay for the detection of feline lentivirus-reactive antibodies in wild felids, employing a puma lentivirus-derived synthetic peptide antigen—M. VAN VUUREN, E. STYLIANIDES (nèe DE KLERK), S.A. KANIA, E.E. ZUCKERMAN and W.D. HARDY Jr 1–6

Rabies

Seroconversion in captive African wild dogs (*Lycaon pictus*) following administration of a chicken head bait/SAG-2 oral rabies vaccine combination—D.L. KNOBEL, A. LIEBENBERG and J.T. DU TOIT 73–77

Reverse-transcription polymerase chain reaction (RT-PCR)

Molecular analysis of bovine viral diarrhoea virus isolates from South Africa—N. KABONGO, C. BAULE and M. VAN VUUREN 273–279

Rhipicephalus gertrudae

Parasites of domestic and wild animals in South Africa. XLIII. Ixodid ticks of domestic dogs and cats in the Western Cape Province—I.G. HORAK and SONJA MATTHEE 187–195

Rhipicephalus sanguineus

Parasites of domestic and wild animals in South Africa. XLIII. Ixodid ticks of domestic dogs and cats in the Western Cape Province—I.G. HORAK and SONJA MATTHEE 187–195

First evidence of *Anaplasma platys* in *Rhipicephalus sanguineus* (Acari: Ixodida) collected from dogs in Africa—Y.O. SANOGO, B. DAVOUST, H. INOKUMA, J-L. CAMICAS, P. PAROLA and P. BROUQUI 205–212

Rickettsaemia

Comparing the detection of exposure to *Ehrlichia ruminantium* infection on a heartwater-endemic farm by the pCS20 polymerase chain reaction assay and an indirect MAP1-B enzyme linked immunosorbent assay—B.H. SIMBI, T.F. PETER, M.J. BURRIDGE and S.M. MAHAN 231–235

Rift Valley fever virus

Indirect enzyme-linked immunosorbent assay for the detection of antibody against Rift Valley fever virus in domestic and wild ruminant sera—J.T. PAWESKA, S.J. SMITH, I.M. WRIGHT, R. WILLIAMS, A.S. COHEN, A.A. VAN DIJK, A.A. GROBBELAAR, J.E. CROFT, R. SWANEPOEL and G.H. GERDES 49–64

Schistosoma mattheei

Schistosoma mattheei—an ovum containing twin miracidia—L.J. VAN RENSBURG and J.A. VAN WYK 69–71

Description and classification of abnormal cercariae of *Schistosoma mattheei* and a method of concentration and collection—J.A. VAN WYK and L.J. VAN RENSBURG 121–129

Scrub hares

Ixodid ticks of Angora and Boer goats, grysbok, common duikers, kudus and scrub hares in Valley Bushveld in the Eastern Cape Province, South Africa—K.M. DE F. MACIVOR and I.G. HORAK 113–120

Setaria saegeri

Studies on the genus *Setaria* Viborg, 1795 in South Africa. II. *Setaria scalprum* (Von Linstow, 1908) and *Setaria saegeri* (Le Van Hoa, 1961)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 7–13

Setaria scalprum

Studies on the genus *Setaria* Viborg, 1795 in South Africa. II. *Setaria scalprum* (Von Linstow, 1908) and *Setaria saegeri* (Le Van Hoa, 1961)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 7–13

Sheep

Ventriculo-atrial conduction in the ovine heart, caused by premature ventricular complexes—J. KER and E.C. WEBB 107–111

Parasites of domestic and wild animals in South Africa. XLII. Helminths of sheep on four farms in the Eastern Cape Province—I.G. HORAK 175–186

The heart remembers: observations of cardiac memory in the Dorper sheep heart—J. KER, E.C. WEBB, J.A. KER and P.A. BEKKER 299–305

Simulium

Review of blackfly (Diptera: Simuliidae) control in South Africa—E. MYBURGH and E.M. NEVILL 307–316

Siphonaptera, Chimaeropsyllidae

Fleas of elephant shrews (Mammalia, Macroscelididae), and a new host and locality record for *Macroscelidopsylla albertyni* De Meillon & Marcus, 1958 (Siphonaptera, Chimaeropsyllidae)—J-C. BEAUCOURNU, I.G. HORAK and L.J. FOURIE 251–253

Sleeping sickness

Occurrence, clinical manifestation and the epidemiological implications of naturally occurring canine trypanosomosis in western Kenya—G. MATETE 317–323

South Africa

Studies on the genus *Setaria* Viborg, 1795 in South Africa. II. *Setaria scalprum* (Von Linstow, 1908) and *Setaria saegeri* (Le Van Hoa, 1961)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 7–13

Parasites of South African wildlife. XVII. *Ostertagia triquetra* n. sp. (Nematoda: Trichostrongylina) from the grey rhebuck, *Pelea capreolus* (Forster, 1790)—J. BOOMKER and M-C. DURETTE-DESSET 37–41

Climate change decreases habitat suitability for some tick species (Acari: Ixodidae) in South Africa—A. ESTRADA-PEÑA 79–93

Clinical and pathological observations on *Streptococcus* sp. infection on South African trout farms with gas supersaturated water supplies—K.D.A. HUCHZERMEYER 95–105

Ixodid ticks of Angora and Boer goats, grysbok, common duikers, kudus and scrub hares in Valley Bushveld in the Eastern Cape Province, South Africa—K.M. DE F. MACIVOR and I.G. HORAK 113–120

Parasites of domestic and wild animals in South Africa. XLI. Arthropod parasites of impalas, *Aepyceros melampus* in the Kruger National Park—I.G. HORAK, G.J. GALLIVAN, L.E.O. BRAACK, J. BOOMKER and V. DE VOS 131–163

Parasites of domestic and wild animals in South Africa. XLII. Helminths of sheep on four farms in the Eastern Cape Province—I.G. HORAK 175–186

Parasites of domestic and wild animals in South Africa. XLIII. Ixodid ticks of domestic dogs and cats in the Western Cape Province—I.G. HORAK and SONJA MATTHEE 187–195

Molecular analysis of bovine viral diarrhoea virus isolates from South Africa—N. KABONGO, C. BAULE and M. VAN VUUREN 273–279

Review of blackfly (Diptera: Simuliidae) control in South Africa—E. MYBURGH and E.M. NEVILL 307–316

Population structure of *Argas arboreus* (Acari: Argasidae) ticks associated with seasonally abandoned mixed heronries, dominated by cattle egrets (*Bubulcus ibis*), in South Africa—V.N. BELOZEROV, D.J. VAN NIEKERK and H.J.B. BUTLER 325–330

***Streptococcus* sp.**

Clinical and pathological observations on *Streptococcus* sp. infection on South African trout farms with gas supersaturated water supplies—K.D.A. HUCHZERMEYER 95–105

Th1 type immune response

Leishmania donovani-derived lipophosphoglycan plus BCG induces a Th1 type immune response but does not protect Syrian golden hamsters (*Mesocricetus auratus*) and BALB/c mice against *Leishmania donovani*—W.K. TONUI, S.S. MPOKE, A.S. ORAGO, S.J. TURCO, P.A. MBATI and G.M. MKOJI 255–263

Ticks

Climate change decreases habitat suitability for some tick species (Acari: Ixodidae) in South Africa—A. ESTRADA-PEÑA 79–93

Ixodid ticks of Angora and Boer goats, grysbok, common duikers, kudus and scrub hares in Valley Bushveld in the Eastern Cape Province, South Africa—K.M. DE F. MACIVOR and I.G. HORAK 113–120

Comparing the detection of exposure to *Ehrlichia ruminantium* infection on a heartwater-endemic farm by the pCS20 polymerase chain reaction assay and an indirect MAP1-B enzyme linked immunosorbent assay—B.H. SIMBI, T.F. PETER, M.J. BURRIDGE and S.M. MAHAN 231–235

Population structure of *Argas arboreus* (Acari: Argasidae) ticks associated with seasonally abandoned mixed heronries, dominated by cattle egrets (*Bubulcus ibis*), in South Africa—V.N. BELOZEROV, D.J. VAN NIEKERK and H.J.B. BUTLER 325–330

Toxoplasma gondii

Isolation of *Toxoplasma gondii* from goats with a history of reproductive disorders and the prevalence of *Toxoplasma* and chlamydial antibodies—S.P. SHARMA, E.K. BAIPOLEDI, J.F.C. NYANGE and L. TLAGAE 65–68

Trichinella zimbabwensis

Effect of host age in the distribution of adult *Trichinella zimbabwensis* in the small intestines of golden hamsters (*Mesocricetus auratus*) and Balb C mice—S. MUKARATIRWA, E. NKULUNGO, ELIZABETH MATENGA and E. BHEBHE 169–173

Trout

Clinical and pathological observations on *Streptococcus* sp. infection on South African trout farms with gas supersaturated water supplies—K.D.A. HUCHZERMEYER 95–105

Trypanocidal drugs

Resistance to trypanocidal drugs—suggestions from field survey on drug use in Kwale district, Kenya—G.L. MUGUNIERI and G.A. MURILLA 29–36

Trypanosomosis

A comparative study on the clinical, parasitological and molecular diagnosis of bovine trypanosomosis in Uganda—J.W. MAGONA, J.S.P. MAYENDE, W. OLAHO-MUKANI, P.G. COLEMAN, N.N. JONSSON, S.C. WELBURN and M.C. EISLER 213–218

Occurrence, clinical manifestation and the epidemiological implications of naturally occurring canine trypanosomosis in western Kenya—G. MATETE 317–323

Tulp poisoning

Conditioned feed aversion as a means to prevent tulp (*Homeria pallida*) poisoning in cattle—L.D. SNYMAN, R.A. SCHULTZ, J.P.J. JOUBERT, K.M. BASSON and L. LABUSCHAGNE 43–48

Uganda

A comparative study on the clinical, parasitological and molecular diagnosis of bovine trypanosomosis in Uganda—J.W. MAGONA, J.S.P. MAYENDE, W. OLAHO-MUKANI, P.G. COLEMAN, N.N. JONSSON, S.C. WELBURN and M.C. EISLER 213–218

Vaccine

Seroconversion in captive African wild dogs (*Lycaon pictus*) following administration of a chicken head bait/SAG-2 oral rabies vaccine combination—D.L. KNOBEL, A. LIEBENBERG and J.T. DU TOIT 73–77

PCR-based detection of the transovarial transmission of Uruguayan *Babesia bovis* and *Babesia bigemina* vaccine strains—V. GAYO, M. ROMITO, L. NEL, M.A. SOLARI and G.J. VILJOEN 197–204

Ventriculo-atrial conduction

Ventriculo-atrial conduction in the ovine heart, caused by premature ventricular complexes—J. KER and E.C. WEBB 107–111

Virus

Evaluation of an indirect enzyme-linked immunosorbent assay for the detection of feline lentivirus-reactive antibodies in wild felids, employing a puma lentivirus-derived synthetic peptide antigen—M. VAN VUUREN, E. STYLIANIDES (nèe DE KLERK), S.A. KANIA, E.E. ZUCKERMAN and W.D. HARDY Jr 1–6

Indirect enzyme-linked immunosorbent assay for the detection of antibody against Rift Valley fever virus in domestic and wild ruminant sera—J.T. PAWESKA, S.J. SMITH, I.M. WRIGHT, R. WILLIAMS, A.S. COHEN, A.A. VAN DIJK, A.A. GROBBELAAR, J.E. CROFT, R. SWANEPOEL and G.H. GERDES 49–64

Molecular analysis of bovine viral diarrhoea virus isolates from South Africa—N. KABONGO, C. BAULE and M. VAN VUUREN 273–279

Western blot

Evaluation of an indirect enzyme-linked immunosorbent assay for the detection of feline lentivirus-reactive antibodies in wild felids, employing a puma lentivirus-derived synthetic peptide antigen—M. VAN VUUREN, E. STYLIANIDES (nèe DE KLERK), S.A. KANIA, E.E. ZUCKERMAN and W.D. HARDY Jr 1–6

Wildlife

Evaluation of an indirect enzyme-linked immunosorbent assay for the detection of feline lentivirus-reactive antibodies in wild felids, employing a puma lentivirus-derived synthetic peptide antigen—M. VAN VUUREN, E. STYLIANIDES (nèe DE KLERK), S.A. KANIA, E.E. ZUCKERMAN and W.D. HARDY Jr 1–6

Studies on the genus *Setaria* Viborg, 1795 in South Africa. II. *Setaria scalprum* (Von Linstow, 1908) and *Setaria saegeri* (Le Van Hoa, 1961)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 7–13

The morphological characteristics of the antebrachiocarpal joint of the cheetah (*Acinonyx jubatus*)—L.O.C. OHALE and H.B. GROENEWALD 15–20

Parasites of South African wildlife. XVII. *Ostertagia triquetra* n. sp. (Nematoda: Trichostrongylina) from the grey rhebuck, *Pelea capreolus* (Forster, 1790)—J. BOOMKER and M-C. DURETTE-DESSET 37–41

Indirect enzyme-linked immunosorbent assay for the detection of antibody against Rift Valley fever virus in domestic and wild ruminant sera—J.T. PAWESKA, S.J. SMITH, I.M. WRIGHT, R. WILLIAMS, A.S. COHEN, A.A. VAN DIJK, A.A. GROBBELAAR, J.E. CROFT, R. SWANEPOEL and G.H. GERDES 49–64

Seroconversion in captive African wild dogs (*Lycaon pictus*) following administration of a chicken head bait/SAG-2 oral rabies vaccine combination—D.L. KNOBEL, A. LIEBENBERG and J.T. DU TOIT 73–77

Ixodid ticks of Angora and Boer goats, grysbok, common duikers, kudus and scrub hares in Valley Bushveld in the Eastern Cape Province, South Africa—K.M. DE F. MACIVOR and I.G. HORAK 113–120

Parasites of domestic and wild animals in South Africa. XLI. Arthropod parasites of impalas, *Aepyceros melampus* in the Kruger National Park—I.G. HORAK, G.J. GALLIVAN, L.E.O. BRAACK, J. BOOMKER and V. DE VOS 131–163

Fleas of elephant shrews (Mammalia, Macroscelididae), and a new host and locality record for *Macroscelidopsylla albertyni* De Meillon & Marcus, 1958 (Siphonaptera, Chimaeropsyllidae)—J-C. BEAUCOURNU, I.G. HORAK and L.J. FOURIE 251–253

Zimbabwe

Effect of host age in the distribution of adult *Trichinella zimbabwensis* in the small intestines of golden hamsters (*Mesocricetus auratus*) and Balb C mice—S. MUKARATIRWA, E. NKULUNGO, ELIZABETH MATENGA and E. BHEBHE 169–173

Detection of antibodies to the *Ehrlichia ruminantium* MAP1-B antigen in goat sera from three communal land areas of Zimbabwe by an indirect enzyme-linked immunosorbent assay—O. KAKONO, T. HOVE, D. GEYSEN and S. MAHAN 243–249